

Opinion

The Impact of Climate Change on Fertility

Benjamin S. Walsh,^{1,5} Steven R. Parratt,^{1,5} Ary A. Hoffmann,² David Atkinson,¹ Rhonda R. Snook,³ Amanda Bretman,⁴ and Tom A.R. Price^{1,*}

Rising global temperatures are threatening biodiversity. Studies on the impact of temperature on natural populations usually use lethal or viability thresholds, termed the ‘critical thermal limit’ (CTL). However, this overlooks important sublethal impacts of temperature that could affect species’ persistence. Here we discuss a critical but overlooked trait: fertility, which can deteriorate at temperatures less severe than an organism’s lethal limit. We argue that studies examining the ecological and evolutionary impacts of climate change should consider the ‘thermal fertility limit’ (TFL) of species; we propose that a framework for the design of TFL studies across taxa be developed. Given the importance of fertility for population persistence, understanding how climate change affects TFLs is vital for the assessment of future biodiversity impacts.

Biodiversity Under Climate Change

Climate change will continue to have an increasingly dramatic effect on the global thermal environment [4], including increases in average local temperatures and the frequency of heatwaves [5,6]. These shifts present a major threat to biodiversity and are starting to have severe impacts on the distribution and abundance of natural populations and species [7,8]. The capacity of species to respond ecologically and evolutionarily to the challenges of global thermal change will affect future biodiversity. Determining key thermally sensitive traits across species, and quantification of the ability of species to buffer the effects of thermal stress on these traits, is therefore a critical research priority [9].

Understanding the long-term impacts of climate change on populations requires robust predictive models that can project responses to both current global temperatures and future climate change scenarios. Currently, many such models are based on empirically derived **CTL** (see [Glossary](#)) estimates, which describe the upper and lower temperature bounds beyond which critical biological functions (e.g., movement, respiration) fail [8,10]. Comparative studies have shown that measures of such viability limits more robustly predict the current distributions of many species than measures derived from changes in mean fitness traits under thermal stress [11]. For this reason, CTLs have also been used to infer species’ sensitivity to climate change [8,12–14]. However, using only thermal limits to viability may be misleading because different measures of CTLs do not always correlate within a single species or population, leading to inconsistent estimates of population persistence [15]. It has been suggested that a multitrait approach to thermal tolerance may give more robust estimates of species’ responses to climate change [15]. In particular, the focus of thermal limits needs to move away from the incapacitating and lethal effects of thermal stress, to investigate how sublethal temperatures impact fitness-related traits such as reproduction that are critical for population stability and persistence.

Sensitivity of Fertility to Temperature

Fertility is a major component of individual fitness and is a central determinant of population growth and persistence. Evidence from a wide variety of taxa suggest that the germ line and

Highlights

Extreme temperatures can severely impact fertility in animals, plants, and fungi.

Climate change may render populations extinct due to fertility losses.

To understand these risks, we need a cross-species measure of thermal fertility loss.

We propose researchers investigate the ‘thermal fertility limit’ (TFL) of organisms.

Standardised measures of TFLs may help to predict species at risk under climate change.

¹Institute of Integrative Biology, University of Liverpool, Liverpool, UK

²School of BioSciences, Bio21 Institute, University of Melbourne, Australia

³Department of Zoology, Stockholm University, Stockholm, Sweden

⁴Faculty of Biological Sciences, University of Leeds, Leeds, UK

⁵Authors contributed equally

*Correspondence: T.Price@liverpool.ac.uk (T.A.R. Price).

associated reproductive physiology is sensitive to thermal stress, particularly high temperatures [16–20]. Evidence, mostly from pollen development, suggests that meiosis is a more thermally sensitive process than mitosis (reviewed in [21,22]). In mammals, the descended testicle has evolved to ensure that spermatogenesis occurs at cooler-than-body temperatures ([23] and references therein). Temperature-induced infertility imposes major economic costs in tropical climates [24]. However, although a number of studies have examined how temperature impacts reproductive traits (Table 1), these often use vastly different methodologies and measure different aspects of reproductive biology. This collection of disparate studies makes quantitative comparisons of the impact of high temperature on reproduction very difficult. Possibly for this reason, thermal limits to fertility have not been systematically incorporated into predictions of species' responses to climate change.

Here we argue that the effect of temperature on fertility requires a broad analogue of CTL, termed the **'Thermal Fertility Limit' (TFL)**. This term would capture both the upper (TF_{MAX}) and lower (TF_{MIN}) temperature boundaries at which a species loses fertility. This new term will facilitate researchers in bringing together related work on how environmental stress impacts this broadly important component of biology and will highlight the important biological and ecological distinction between fertility and survival when assessing species' response to climate change. We suggest that a framework be developed that will allow researchers to design and conduct thermal fertility studies in a way that generates comparable datasets across taxa. A large database of TFL measures across multiple species and populations, relevant to thermal stress levels encountered in nature, would provide the power to answer important evolutionary and ecological questions regarding the impact of climate change on natural populations at risk (Box 1 and Figure I in Box 1). We do not propose that TFL measures would replace CTLs. Rather, we suggest that the combination of these measures, the geographic distribution of these two limits, and the extent to which they correlate within and among species will give valuable insight into species' ability to persist and adapt to global thermal change. To do this, we need to consider how temperature is likely to affect fertility at a mechanistic level and how researchers can design and conduct studies of TFLs in a standardised and broadly comparable way.

Towards a Methodological Framework for the Study of TFLs

The adoption of standardised measures for CTLs [11,25], typically either a direct or a proxy measure of viability, has facilitated large-scale comparative studies of species' responses to climate change [8]. A challenge for the study of TFLs will be to develop a similarly standardised measure for fertility. This is a nontrivial task given the inherent complexity and potential species specificity of reproductive components that contribute to fertility (Figure 1). This complexity is highlighted by the diverse methodologies and metrics of fertility employed in the existing literature on the effect of temperature on fertility (Table 1). For maximum utility, TFL studies should be carefully designed either to produce a quantitative point estimate of temperature limits for fertility for comparative species distribution modelling, or to generate effect size estimates for fertility loss at a given thermal stress level for future meta-analyses between groups.

Factors in Designing TFL Studies

Despite the diverse elements of fertility described in Figure 1, we argue that the most ecologically precise limit to fertility is the point at which the qualitative ability of an organism to produce viable adult offspring under controlled conditions is lost. This limit yields a precise metric that can be applied to quantitative comparisons among taxa. However, for many species measuring offspring production directly may be impractical; for example, if generation times are extremely slow. In such instances, proxy measurements that can be empirically correlated with

Glossary

Critical thermal limits (CTLs): a suite of commonly used measures of the maximum and minimum temperatures at which organisms can viably function. Individuals are exposed to either static stressful temperatures or gradually ramping temperatures and observed for physiological failure (e.g., uncoordinated movement, heat coma, death [1]). Typically, either the duration of exposure or the temperature at which loss of viability is observed is recorded as the thermal limit.

Fecundity: the total number of offspring an individual can produce across a set interval or lifetime.

Fertility: the ability of an organism to produce viable offspring. Fertility can be measured in a number of ways but always reaches its lower limit when conditions prevent an individual from producing any offspring (i.e., sterility).

Hardening: increased thermal tolerance shown by organisms after a short period of exposure to a stressful but nonlethal temperature within the same life stage. Hardening tests are one component of a species' plastic response when exposed to stressful temperatures [2].

Sterility: describes an individual that cannot produce any offspring over a defined period and thus is synonymous with complete infertility.

Thermal fertility limits (TFLs): outlined here for the first time, TFLs refer to a level and duration of thermal stress that renders individuals unable to reproduce. For populations and species, this can be defined as the temperature at which a given proportion of individuals are qualitatively sterile, and it includes both higher (TF_{MAX}) and lower (TF_{MIN}) thermal stress limits. For example, the upper TF_{MAX} of male *Drosophila buzzatii* – measured as permanent sterility of 80% of individuals after 6 h – is 38.5°C [3].

Table 1. Examples of Thermal Impacts on Fertility

Taxonomic group	Organism	Species	Impact of temperature on fertility	Measure	Refs
Cnidarian	Coral	<i>Acropora digitifera</i>	Increase of 2°C reduced the number of sperm bundles by almost 50% and reduced egg size	Gamete number	[49]
Insect	Bed bug	<i>Cimex lectularius</i>	Egg production and hatching success can fall to almost zero as a result of thermal stress	Fecundity	[30]
	Red mason bee	<i>Osmia bicornis</i>	Changed odour profile, altering female mating preference	Mating preference	[50]
	Beetle	<i>Callosobruchus maculatus</i>	Males reared at extreme high temperatures produce smaller sperm than benign controls	Sperm form and function	[51]
	Beetle	<i>Tribolium castaneum</i>	Stressed males reduce sperm viability and competitiveness; inseminated sperm in female storage organs less viable when female stressed; transgenerational impact reducing longevity of offspring sired by stressed males	Sperm form and function, offspring production	[36]
	Dragonfly	<i>Micrathyrja</i> spp.	Species in the genus that struggle to maintain optimal body temperatures are less efficient at defending perches at high temperatures and lose out on breeding sites to larger species	Courtship behaviour	[52]
	Fruit fly	<i>Bactrocera tryoni</i>	Reduced mating latency at cold temperatures, reduced mating frequency at cold temperatures	Mating latency, mating frequency	[53]
	Fruit fly	Family: Drosophilidae	Reduced mating success; impairment of sperm elongation, resulting in loss of sperm motility and thus lower fertility	Offspring production, mating success, sperm motility	[17,26,29,31,54–56]
	Oriental fruit moth	<i>Grapholita molesta</i>	A 2-h heat stress during pupation reduced fecundity but increased other adult fitness traits such as survival	Fecundity, gamete viability	[57]
	Wasp	<i>Aphidius avenae</i>	Low mating success rate due to reduced courtship behaviour; reduced sperm count after developmental stress, with males at high stress fully sterile; reduced fertilisation results in fewer females, secondarily altering sex ratios; stressed females produce fewer eggs	Courtship behaviour, gamete number, fertilisation success, and offspring production	[32,58]
Poales	Barley	<i>Hordeum vulgare</i>	Developing anther cells are compromised during thermal stress while developing ovule cells are not	Gamete viability	[59]
	Rice	<i>Oryza sativa</i>	High temperature during flowering increased pollen sterility, with greater sterility if CO ₂ levels were high	Gamete viability	[60]
Polemoniales	Tomato	<i>Solanum lycopersicum</i>	Under thermal stress pollen viability was reduced and anthers developed abnormalities; thermally tolerant genotypes showed resistance	Gamete viability	[61]
Vertebrate	Chicken	<i>Gallus gallus domesticus</i>	An 8-week thermal stress results in increased sperm death and associated drop in fertility	Sperm concentration	[18]
	Cow	<i>Bos taurus</i>	Ovulation failure and abortion rate is higher in cows inseminated during warm seasons	Fertilisation	[62]
	Guppy fish	<i>Poecilia reticulata</i>	Males raised at stressful temperatures have shorter, slower sperm than individuals raised at benign temperatures	Sperm form and function	[63]
	Mouse	<i>Mus musculus</i>		Gamete number	[16]

Table 1. (continued)

Taxonomic group	Organism	Species	Impact of temperature on fertility	Measure	Refs
			Reduced sperm count for over 60 days after 30-min heat shock		
	Pig	<i>Sus</i> sp.	Sperm DNA damage higher and sperm concentration lower during warm wet season	Sperm form and function	[24]
	Sea lion	<i>Otaria flavescens</i>	Stressed males desert females to thermoregulate, foregoing mating opportunities	Courtship and mating behaviour	[64]
	Zebra finch	<i>Taeniopygia guttata</i>	Daily heatwaves reduced the proportion of sperm exhibiting normal morphology	Sperm form and function	[65]

fertility may also serve to capture the effect of temperature. For example, in some *Drosophila* qualitative sperm motility has been used to quantify male fertility following heat stress, as this correlates strongly with reproductive output (reviewed in [26]). In plants, the percentage of pollen grains that germinate *in vitro* correlates with fruit productivity and has been employed as a measure of TFLs [21,27]. It would be unrealistic to attempt to identify a trait that captures the effect of temperature on fertility across all of biology, but taxon-specific proxies like these may be sufficient to enable meaningful comparative studies.

Whichever measurement is used, assessing fertility over a range of static temperatures will allow us to generate a fertility reaction norm. From these reaction norms we can determine the temperature at which fertility drops by a given percentage compared with benign controls; a measure analogous to a 'lethal dosage' in toxicology and one already used for some measures of CTLs [28]. The exact proportion of fertility loss that is ecologically relevant for population stability and thus represents a true TFL is likely to vary from species to species. With enough data on the reproductive and population biology of a given organism, these thresholds could be

Box 1. Groups at Risk

Examples of Organisms That May Be Particularly at Risk of Losing Fertility Due to High Temperatures

Certain groups of organisms are likely to be most vulnerable to temperature-driven fertility loss. These groups may provide important case studies and primary avenues of research (Figure 1).

Ectothermic Species

Most plant species cannot regulate the temperature of their tissues (excluding a number of species of flower [42]), forcing them to withstand ambient temperatures. Likewise, ectothermic animals may also be vulnerable [5], as they rely on behavioural rather than physiological thermoregulation to avoid stressful microenvironments. Smaller ectothermic animals are even more at risk, as they will reach ambient temperatures faster.

Endemic Species and Species with Small Ranges

Rare or endemic species with small latitudinal ranges are likely to be particularly at risk of losing fertility as ambient temperatures increase because: (i) they are likely to lack the genetic variation and gene flow required to adapt to novel stressors [7]; and (ii) in many cases they may be unable to shift their distribution range to track changing climates. This will be particularly true for island endemics and species that live in specialised elevational niches in mountains.

Aquatic Species

Aquatic species, particularly broadcast spawners, are likely to be at risk because the specific heat capacity of water will result in rapid changes in tissue temperatures. Further, gametes in the water from spawning organisms will be exposed directly to stressful temperatures, so will need to evolve robust physiological responses to high temperatures to retain form and function. This is likely to be a greater issue for freshwater and shallow-water organisms, as these environments experience greater fluctuations in temperatures, exposing these organisms to acute stress events.

Sessile Species and Life Stages

Sessile organisms, such as plants, corals, and juvenile stages (e.g., pupal stages in holometabolous insects), in which movement to cooler areas during temperature spikes is not possible, may be particularly vulnerable. Similarly, due to their limited dispersal ability, belowground communities may be especially vulnerable to fertility loss under climate change [43].




Trends in Ecology & Evolution

Figure 1. Examples of Organisms That May Be Particularly at Risk of Losing Fertility Due to High Temperatures. Clockwise from top left: broadcast-spawning fish such as carp; small ectothermic insects including pollinating bees; endemic animals with limited latitudinal or elevation ranges such as the flightless cormorant; disease vectors including mosquitos; coral species that are important to highly diverse reefs; and endemic plant species including the Scottish primrose. All photographs in this figure are licensed under CC BY 2.0. Credits: Joaquim Alves Gaspar, Charles Sharp, Toby Hudson, and David Glass.

explicitly modelled. Alternatively, if reaction norms are established across a broad enough range of temperatures, it should be possible to determine any threshold and to assess whether these are correlated across species.

Further, unlike viability limits, fertility is not necessarily an irreversible binary trait. Evidence suggests that complete **sterility** at extreme temperatures is preceded by quantitative fertility loss under intermediate conditions [29,30]. Furthermore, recovery of fertility can occur in some heat-sterilised animals if they are returned to benign conditions [31,32], although under severe thermal stress sterility can be permanent ([3,19] and personal observation). Researchers should carefully consider the timeframe over which qualitative fertility is assessed following heat stress and potentially account for the recovery of fertility over time; a 2-day knockdown in fertility may be inconsequential for long-lived species but catastrophic for organisms that exist as adults for only days. This highlights an important consideration when comparing the utility of CTLs and TFLs, reinforcing that TFLs have a much more complicated relationship with time than CTLs.

A second important practical consideration arises when selecting an ecologically relevant temperature treatment. Researchers have shown that the response of organisms to thermal



Trends in Ecology & Evolution

Figure 1. A Generalised and Simplified Schematic of the Stages in Sexual Reproduction and Examples of Organisms for Which the Effect of Temperature Has Been Measured at These Stages (Table 1). Fertility is the emergent product of multiple physiological, developmental, and behavioural processes. Not all steps are relevant to all organisms; indeed, the diversity and complexity of this cascade across sexual organisms is not fully captured here. However, in all cases the 'success' of fertility begins with the generation of gametes and ends with the production of viable offspring. High temperature may perturbate single or multiple

(Figure legend continued on the bottom of the next page.)

stress is affected by both the intensity of the temperature chosen and the duration of exposure [25]. This is further complicated when one considers the effects that **hardening** treatments [1], ramping [13], and the observed differences between static and cyclic temperature treatments ([33] and references therein) have on thermal performance in many organisms. Unlike CTLs, where the effect of temperature is often immediately visible, loss of fertility requires subsequent assays following exposure to heat, and so ramping assays are unlikely to be useful. Instead, researchers must choose regimes of static or fluctuating temperature stress that reflect current or future thermal extremes for natural populations. The need to finely balance high-throughput, standardised, repeatable assays with ecological realism will be a major challenge for TFL research.

To summarise, if researchers think about the exact trait they are going to measure and the thermal regime under which it will be measured, and consider that fertility may recover over time, they will be well on their way to having a robust framework for the study of TFLs (Box 2). Investigating this in model species, and testing whether it predicts species distributions better than current methods, will be a key step in determining how important TFLs are in nature.

Can Species Maintain Fertility in the Face of Thermal Change?

Many species are predicted to have populations pushed beyond their critical thermal maxima (CT_{MAX}) by climate change [14]. As thermal fertility maxima (TF_{MAX}) are expected to often be lower than CT_{MAX} , rapid climate change is likely to push many populations and species beyond their TF_{MAX} . The development of standardised measures of TFLs will provide tools to investigate how species might physiologically acclimate and adapt to these changing thermal environments.

Are TFLs Plastic?

Organisms could show phenotypic plasticity in TFLs within their own lifetime or through intergenerational carry-over effects. Suboptimal temperatures experienced at early life-history stages can affect traits such as adult size [34]. Experiencing some level of thermal stress can increase the fitness of individuals for a similar stress later in life, a process known as acclimation. For CTLs there is significant, but very limited, scope for coping with rising temperatures through plasticity [35]. For instance, the degree of plasticity in upper thermal tolerance appears weakly associated with species' distribution ranges [13]. However, it is not known whether similar plasticity exists for TFLs or whether plasticity in TFLs is greater than that for CTLs. Exposing organisms to acclimation treatments followed by TFL measurement, or investigating intergeneration carry-over effects for TFLs, may shed new light on the ability of organisms to buffer the effects on fitness of ecological change.

There is mixed evidence for the impact of acclimation on temperature-induced sterility. Male *Drosophila buzzatii* regain fertility faster following a heat stress if they had previously experienced a heat shock [3]. However, both *Drosophila subobscura* and *Tribolium castaneum* have been shown to exhibit more extreme fertility loss when exposed to multiple rather than single periods of heat stress, which does not indicate an acclimation response [17,36]. Where plasticity in thermal fertility traits does exist, the underlying mechanisms remain largely unknown. However, individuals are likely to cope with stress in part by using heat-shock proteins, which are important in mediating upper thermal limits in insect species [37]. Many, including Hsp70,

steps in this process, but early meiotic stages can be particularly thermally sensitive [21]. High temperature may affect several of these traits simultaneously in an individual; for example, by both arresting gametogenesis and reducing investment in copulation behaviours. By contrast, the effect of high temperature on a single trait, say testis development, may subsequently have cascading effects on downstream elements of reproduction such as sperm count and motility. Photograph credits: (A) barley, Raul Dupagne; (B) guppy, Baskua; (C) *Drosophila* mating, D. Chai; (D) coral reef, Toby Hudson; (E) rooster, Pete Linforth. All photographs licensed under CC BY 2.0.

Box 2. Considerations When Designing TFL Experiments

Trait Selection

We suggest that wherever possible researchers measure both qualitative and quantitative offspring production to capture the ecological impact of high temperature on fertility. Where this is impossible, careful selection of proxy measures of fertility that can be empirically correlated with an individual's ability to produce offspring could be considered. Holistic measures such as these are most likely to generate broadly comparable data sets across taxa.

Life-History Stage

While reproduction occurs almost invariably during adult life-history stages, reproductive development and maturation can begin much earlier. Researchers should therefore consider which life stage of their organism to expose to stress. For instance, do heat-treated juveniles mature into sterile adults while heated adults remain fertile?

Ecologically Valid Thermal Environment

Careful attention should be given to selecting temperature regimes that reflect the current or future extremes that organisms are likely to face. For instance, are temperature spikes over a matter of a few hours more likely to impact a species' fertility than a rise in mean daytime temperature? A large body of work on CTLs has demonstrated that measures of thermal performance can be highly sensitive to the duration of stress [25], rates of temperature ramping [13], and the intensity and frequency of any temperature fluctuations [44]. The latter point in particular may be key for thermal fertility, as some animals can recover fertility during periods of benign temperatures including night time [45]. Once researchers have selected a regime of temperature delivery they should strive, where possible, to measure thermal fertility over a range of temperature values. This will help to capture the thermal fertility reaction norm of their organism.

Implications for Population Stability

To estimate the population-level effects of high temperature on fertility, researchers should consider what percentage loss of fertility represents a meaningful threat to population stability. Factors such as the effective population size of the organism in nature, the potential **fecundity** of individuals, and their generation time could be used to estimate a species' sensitivity to fertility loss. Researchers can then determine the degree of thermal stress required to push their study organism beyond this threshold.

Critical Thermal and Fertility Limits

The power of TFLs to predict species' response to climate change will be related to the extent to which fertility and viability limits correlate with each other and across species. Low correlation would suggest that one metric cannot be substituted for the other. Which species have high and which species have low correlation and what impacts this relationship? Thus, researchers should determine both the fertility and the viability limits of their organism under relevant thermal regimes.

are upregulated during hardening treatments, helping individuals to offset the negative fitness consequences of thermal stress [38]. Heat-shock proteins are a ubiquitous component in living systems; importantly, they are found in gametes, including human spermatozoa [39]. Exploring the scope for heat-shock protein expression to buffer the deleterious effect of high temperature on fertility and the variation in this within closely related species might explain patterns of variation in TFLs.

Can TFLs Evolve?

Over long periods of environmental change, selection should favour more thermally tolerant genotypes and a rise in both CTLs and TFLs. Including the evolvability of thermally sensitive traits in models of species' response to climate change generates vastly different predictions than equivalent models parameterised with only current measures of thermal sensitivity [8]. However, current evidence suggests there is very little standing genetic variation and evolvability for high-temperature CTLs [8], although this is debated (reviewed in [25]). Whether TFLs can evolve rapidly is unknown. Limited evidence in *Drosophila* has shown that male sterility under heat stress can be variable within species and may be under selection to be locally adapted across populations originating from different thermal regimes [17,19,31,40], suggesting that TFLs may be evolvable. Quantification of standing variation in TFLs across genotypes and populations of multiple species would be a good first approach to test this.

Species with CTLs that are low and evolutionarily constrained are predicted to be at particular risk from climate change [12]. For instance, tropical species have been shown to often lack genetic variation that would enable rapid evolution to cope with changing climatic variables such as temperature and desiccation [14,41]. Establishing how these species' TFLs respond to increasing temperatures may be critical to predict how they will be impacted by climate change. If TFLs are substantially lower than CTLs, these species may be more vulnerable than currently predicted. However, if TFLs are more evolvable than CTLs, this may compensate for their initially low TFLs, making CTLs more important predictors of distributions in a warming world. Until both CTLs and TFLs are examined across a variety of taxa, and the evolvability of TFLs determined, confidence in predictions about which taxa are going to be particularly vulnerable will be low (Box 1).

Whether populations or species can respond to thermally induced loss of fertility, through either short-term plasticity or long-term adaptive change, is unclear. This is partly because of knowledge gaps regarding the impact of extreme temperature on fertility in animals and plants. A fundamental understanding of how extreme increases and decreases in temperature influence reproduction with negative effects on fertility is required before the ecological relevance and potential evolution of TFLs can be determined. However, it is precisely these answers that are ultimately among the most important to know, as they will improve predictions of how climate change may affect species' abundance and distribution and thereby change biodiversity across the globe.

Concluding Remarks

Here we have introduced and discussed the idea that measuring the thermal limit of fertility across multiple species and a broad range of taxa could be critical when assessing the impacts of global thermal change on biodiversity. While the use of critical thermal limits has proved to be informative for modelling current and future distributions of species [8,13,14], CTLs may overestimate species' ability to cope with stressful temperatures. Research exploring TFLs (see Outstanding Questions) is needed to ascertain the extent to which they correlate with CTLs. To this end, we propose a general framework for TFL studies to promote large-scale, cross-taxon assessments of this important but largely neglected trait. Focusing on TFLs with broadly standardised methodologies may improve our knowledge of how climate change will affect species' abundance, distribution, and persistence. However, the current literature on how thermal stress impacts fertility is fragmented. Stronger and more unified thermal fertility research might radically improve our predictions about the impacts of global thermal change.

Acknowledgments

The authors thank the Editor and three anonymous reviewers for their constructive comments on this Opinion article. This work was funded by the UK Natural Environment Research Council (grant NE/P002692/1 to T.A.R.P., A.B., and R.R.S.).

References

1. Overgaard, J. *et al.* (2012) Validity of thermal ramping assays used to assess thermal tolerance in arthropods. *PLoS One* 7, e32758
2. Loeschcke, V. and Hoffmann, A.A. (2007) Consequences of heat hardening on a field fitness component in *Drosophila* depend on environmental temperature. *Am. Nat.* 169, 175–183
3. Jørgensen, K.T. *et al.* (2006) Heat tolerance and the effect of mild heat stress on reproductive characters in *Drosophila buzzatii* males. *J. Therm. Biol.* 31, 280–286
4. Intergovernmental Panel on Climate Change (2014) *Climate Change 2014—Impacts, Adaptation and Vulnerability: Regional Aspects*, Cambridge University Press
5. Kingsolver, J.G. *et al.* (2013) Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Funct. Ecol.* 27, 1415–1423
6. Buckley, L.B. and Huey, R.B. (2016) Temperature extremes: geographic patterns, recent changes, and implications for organismal vulnerabilities. *Glob. Chang. Biol.* 22, 3829–3842
7. Hoffmann, A. (2010) Physiological climatic limits in *Drosophila*: patterns and implications. *J. Exp. Biol.* 213, 870–880
8. Kellermann, V. *et al.* (2012) Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. *Proc. Natl Acad. Sci. U. S. A.* 109, 16228–16233

Outstanding Questions

Species Distributions

A major question that will arise as TFLs are investigated is: how well do current species distributions map to a species' TFL and how does this differ from its CTL? Also, how do future species distributions, using the IPCC's predictions, change when modelling TFL against CTL? We already know that different species' average CTL can predict current distributions effectively [8,12,46], although the literature here is largely confined to arthropods. Adding data on TFLs to these model taxa would be an important first step in testing whether TFLs are useful.

There are two ways in which a species' TFL could be more useful than CTL when mapping onto current and future species distributions. First, a species' TFL is likely to be lower than its CTL, which should provide a more accurate measure for critical effects of temperatures. Second, if the correlation between TFL and CTL is inconsistent between species, this could change the predictions of distribution models, as it may reveal particularly vulnerable species that were thought to be safe and vice versa.

Another important question is: can TFLs change across a single species' range? There may be clines in TFLs from high to low latitudes. There is currently some evidence for this, mostly in arthropods [17,19,31], but generalisation across species groups will be important to estimate the variation and evolvability of TFLs within species.

Sexual Traits

Sexual traits are among the most variable traits in biology, with closely related species showing considerable differences in mating behaviour, age at sexual maturity, gamete size, and gamete form. Much of this diversity remains unexplained. For example, some species in the family Drosophilidae produce sperm over ten times longer than their body, while related species produce sperm 100 times smaller [47]. Yet the evolutionary benefits of possessing such large sperm are largely unknown. Assessing how temperature may affect gametes,

9. Moritz, C. and Agudo, R. (2013) The future of species under climate change: resilience or decline? *Science* 341, 504–508
10. Geerts, A. *et al.* (2015) Rapid evolution of thermal tolerance in the water flea. *Daphnia*. *Nat. Clim. Chang.* 5, 665–668
11. Overgaard, J. *et al.* (2014) Sensitivity to thermal extremes in Australian *Drosophila* implies similar impacts of climate change on the distribution of widespread and tropical species. *Glob. Chang. Biol.* 20, 1738–1750
12. Bush, A. *et al.* (2016) Incorporating evolutionary adaptation in species distribution modelling reduces projected vulnerability to climate change. *Ecol. Lett.* 19, 1468–1478
13. Mitchell, K.A. *et al.* (2011) Phenotypic plasticity in upper thermal limits is weakly related to *Drosophila* species distributions. *Funct. Ecol.* 25, 661–670
14. Kellermann, V. *et al.* (2009) Fundamental evolutionary limits in ecological traits drive *Drosophila* species distributions. *Science* 325, 1244–1246
15. Blackburn, S. *et al.* (2014) Evolutionary capacity of upper thermal limits: beyond single trait assessments. *J. Exp. Biol.* 217, 1918–1924
16. Pérez-Crespo, M. *et al.* (2008) Scrotal heat stress effects on sperm viability, sperm DNA integrity, and the offspring sex ratio in mice. *Mol. Reprod. Dev.* 75, 40–47
17. Porcelli, D. *et al.* (2016) Local adaptation of reproductive performance during thermal stress. *J. Evol. Biol.* 30, 422–429
18. Karaca, A. *et al.* (2002) The effects of heat stress and sperm quality classification on broiler breeder male fertility and semen ion concentrations. *Br. Poult. Sci.* 43, 621–628
19. Vollmer, J. *et al.* (2004) Heat and cold-induced male sterility in *Drosophila buzzatii*: genetic variation among populations for the duration of sterility. *Heredity* 92, 257–262
20. Reinhardt, K. *et al.* (2015) An ecology of sperm: sperm diversification by natural selection. *Annu. Rev. Ecol. Evol. Syst.* 46, 435–459
21. Sage, T.L. *et al.* (2015) The effect of high temperature stress on male and female reproduction in plants. *Field Crops Res.* 182, 30–42
22. Paupière, M.J. *et al.* (2014) The metabolic basis of pollen thermo-tolerance: perspectives for breeding. *Metabolites* 4, 889–920
23. Moreno, R.D. *et al.* (2012) Molecular basis of heat stress damage in mammalian testis. In *Testis: Anatomy, Physiology and Pathology* (Nemoto, Y. and Inaba, N., eds), pp. 127–155, Nova Science
24. Peña, S.T. *et al.* (2018) Tropical summer induces DNA fragmentation in boar spermatozoa: implications for evaluating seasonal infertility. *Reprod. Fertil. Dev.* Published online November 12, 2018. <http://dx.doi.org/10.1071/RD18159>
25. Terblanche, J.S. *et al.* (2007) Critical thermal limits depend on methodological context. *Proc. Biol. Sci.* 274, 2935–2943
26. David, J. *et al.* (2005) Male sterility at extreme temperatures: a significant but neglected phenomenon for understanding *Drosophila* climatic adaptations. *J. Evol. Biol.* 18, 838–846
27. Acar, I. and Kakani, V.G. (2010) The effects of temperature on *in vitro* pollen germination and pollen tube growth of *Pistacia* spp. *Sci. Hort.* 125, 569–572
28. Lutterschmidt, W.I. and Hutchison, V.H. (1997) The critical thermal maximum: history and critique. *Can. J. Zool.* 75, 1561–1574
29. Chakir, M. *et al.* (2002) Male sterility thermal thresholds in *Drosophila*: *D. simulans* appears more cold-adapted than its sibling *D. melanogaster*. *Genetica* 114, 195–205
30. Rukke, B.A. *et al.* (2018) Temperature stress deteriorates bed bug (*Cimex lectularius*) populations through decreased survival, fecundity and offspring success. *PLoS One* 13, e0193788
31. Rohmer, C. *et al.* (2004) Heat induced male sterility in *Drosophila melanogaster*: adaptive genetic variations among geographic populations and role of the Y chromosome. *J. Exp. Biol.* 207, 2735–2743
32. Nguyen, T.M. *et al.* (2013) Heat stress affects male reproduction in a parasitoid wasp. *J. Insect. Physiol.* 59, 248–254
33. Sgrò, C.M. *et al.* (2016) What can plasticity contribute to insect responses to climate change? *Annu. Rev. Entomol.* 61, 433–451
34. Atkinson, D. (1994) Temperature and organism size: a biological law for ectotherms? *Adv. Ecol. Res.* 25, 1–58
35. Sørensen, J.G. *et al.* (2016) Evolutionary and ecological patterns of thermal acclimation capacity in *Drosophila*: is it important for keeping up with climate change? *Curr. Opin. Insect Sci.* 17, 98–104
36. Sales, K. *et al.* (2018) Experimental heatwaves compromise sperm function and cause transgenerational damage in a model insect. *Nat. Commun.* 9, 4771
37. Krebs, R. and Loeschcke, V. (1994) Costs and benefits of activation of the heat-shock response in *Drosophila melanogaster*. *Funct. Ecol.* 8, 730–737
38. Sørensen, J. *et al.* (2001) Genetic variation in thermal tolerance among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression and variation in heat stress resistance traits. *Funct. Ecol.* 15, 289–296
39. Miller, D. *et al.* (1992) Characterization and cellular distribution of human spermatozoal heat shock proteins. *Hum. Reprod.* 7, 637–645
40. Pedersen, L.D. *et al.* (2011) The effects of inbreeding and heat stress on male sterility in *Drosophila melanogaster*. *Biol. J. Linn. Soc. Lond.* 104, 432–442
41. Deutsch, C.A. *et al.* (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl. Acad. Sci. U. S. A.* 105, 6668–6672
42. Watling, J.R. *et al.* (2008) Mechanisms of thermoregulation in plants. *Plant Signal. Behav.* 3, 595–597
43. Berg, M.P. *et al.* (2010) Adapt or disperse: understanding species persistence in a changing world. *Glob. Chang. Biol.* 16, 587–598
44. Davies, C. *et al.* (2016) Effect of stable and fluctuating temperatures on the life history traits of *Anopheles arabiensis* and *An. quadriannulatus* under conditions of inter- and intra-specific competition. *Parasit. Vectors* 9, 342
45. Zhao, F. *et al.* (2014) Night warming on hot days produces novel impacts on development, survival and reproduction in a small arthropod. *J. Anim. Ecol.* 83, 769–778
46. Hoffmann, A.A. *et al.* (2003) Adaptation of *Drosophila* to temperature extremes: bringing together quantitative and molecular approaches. *J. Therm. Biol.* 28, 175–216
47. Pitnick, S. *et al.* (1995) Delayed male maturity is a cost of producing large sperm in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* 92, 10614–10618
48. Cooper, E.J. (2014) Warmer shorter winters disrupt Arctic terrestrial ecosystems. *Annu. Rev. Ecol. Evol. Syst.* 45, 271–295
49. Paxton, C.W. *et al.* (2016) Effect of elevated temperature on fecundity and reproductive timing in the coral *Acropora digitifera*. *Zygote* 24, 511–516
50. Conrad, T. *et al.* (2017) The effect of temperature on male mating signals and female choice in the red mason bee, *Osmia bicornis* (L.). *Ecol. Evol.* 7, 8966–8975
51. Vasudeva, R. *et al.* (2014) Developmental temperature affects the expression of ejaculatory traits and the outcome of sperm competition in *Callosobruchus maculatus*. *J. Evol. Biol.* 27, 1811–1818
52. May, M.L. (1977) Thermoregulation and reproductive activity in tropical dragonflies of the genus *Micrathyrja*. *Ecology* 58, 787–798
53. Meats, A. and Fay, F. (2000) Distribution of mating frequency among males of the Queensland fruit fly, '*Bactrocera tryoni*' (Froggatt), in relation to temperature, acclimation and chance. *Gen. Appl. Ent.* 29, 27
54. Gefen, E. and Gibbs, A.G. (2009) Interactions between environmental stress and male mating success may enhance evolutionary divergence of stress-resistant *Drosophila* populations. *Evolution* 63, 1653–1659
55. Arapepe, L. *et al.* (2004) Male sterility thresholds in a tropical cosmopolitan drosophilid, *Zaprionus indianus*. *J. Therm. Biol.* 29, 73–80

behaviour, and other components of fertility may help to explain variation in sexual traits similar to this across the animal kingdom.

Upper or Lower TFLs?

While mean temperatures are expected to rise and temperature spikes become more common due to global climate change, some organisms are also likely to face colder conditions than their current habitat, especially if insulating snow coverage is reduced (e.g. [48]). It will therefore be important to establish the thermal fertility window for many species by measuring both the maximum TF (TF_{MAX}) and the minimum TF (TF_{MIN}). Comparisons between minimum and maximum critical limits within species have found that these traits vary in their evolvability; typically, lower limits have proved to be more variable and evolutionarily responsive. Establishing the extent to which TF_{MAX} and TF_{MIN} correlate within species and whether these traits differ in their standing variation or plasticity will be an important future avenue of research.

56. Batista, M.R.D. *et al.* (2018) Altitudinal distribution of two sibling species of the *Drosophila tripunctata* group in a preserved tropical forest and their male sterility thermal thresholds. *J. Therm. Biol.* 71, 69–73
57. Zheng, J. *et al.* (2017) Are adult life history traits in oriental fruit moth affected by a mild pupal heat stress? *J. Insect. Physiol.* 102, 36–41
58. Roux, O. *et al.* (2010) How does heat shock affect the life history traits of adults and progeny of the aphid parasitoid *Aphidius avenae* (Hymenoptera: Aphidiidae)? *Bull. Entomol. Res.* 100, 543–549
59. Oshino, T. *et al.* (2007) Premature progression of anther early developmental programs accompanied by comprehensive alterations in transcription during high-temperature injury in barley plants. *Mol. Genet. Genomics* 278, 31–42
60. Matsui, T. *et al.* (1997) Effects of high temperature and CO₂ concentration on spikelet sterility in indica rice. *Field Crops Res.* 51, 213–219
61. Müller, F. *et al.* (2016) High-temperature-induced defects in tomato (*Solanum lycopersicum*) anther and pollen development are associated with reduced expression of B-class floral patterning genes. *PLoS One* 11, e0167614
62. De Rensis, F. *et al.* (2017) Causes of declining fertility in dairy cows during the warm season. *Theriogenology* 91, 145–153
63. Breckels, R.D. and Neff, B.D. (2013) The effects of elevated temperature on the sexual traits, immunology and survivorship of a tropical ectotherm. *J. Exp. Biol.* 216, 2658–2664
64. Campagna, C. and Le Boeuf, B.J. (1988) Thermoregulatory behaviour of southern sea lions and its effect on mating strategies. *Behaviour* 107, 72–89
65. Hurley, L.L. *et al.* (2018) Experimental heatwaves negatively impact sperm quality in the zebra finch. *Proc. Biol. Sci.* 285, 20172547